

Guide to Talking to Your Doctor

Print out this guide and keep it in a convenient place.

Share it with your family, loved ones, and anyone who's coming with you to the doctor.

Before your next doctor's appointment, go to **MULTAQ.com** and read the MULTAQ® Medication Guide on page 5 of the Prescribing Information. Then use this checklist of questions to guide a conversation with your doctor about MULTAQ.

And if you've already been prescribed MULTAQ, there are some additional important questions you may want to ask your doctor on the following page.

MULTAQ is a prescription medicine used to lower the chance that you would need to go into the hospital for atrial fibrillation (AFib). It is meant for people who have had AFib in the past but are now in normal rhythm.

IMPORTANT UPDATE: Please see the latest Safety Information on heart failure and permanent AFib as well as the updated MULTAQ Medication Guide on **MULTAQ.com**

If you're interested in talking to your doctor about MULTAQ® (dronedaron) 400 mg Tablets:

Question: Is MULTAQ right for me?

Notes: _____

Question: Can MULTAQ help reduce my risk of going to the hospital for AFib?

Notes: _____

Question: How is MULTAQ different from the other medications I'm taking?

Notes: _____

Question: What are the serious risks associated with MULTAQ that I should know about?

Notes: _____

Question: Will MULTAQ interfere with any of the other medications I'm taking?

Notes: _____

Important Safety Information for MULTAQ

Do not take MULTAQ if you have symptoms of heart failure which have recently worsened or if you have severe heart failure. People with these conditions who take MULTAQ have an increased chance of dying.

Do not take MULTAQ if you have a type of atrial fibrillation (AFib) called permanent AFib. You should not take MULTAQ if you are in AFib and your doctor is not planning to change you back into normal rhythm.

Call your doctor immediately if you're taking MULTAQ and have any signs and symptoms of heart failure. These may include shortness of breath or wheezing at rest; wheezing, chest tightness, or coughing up frothy sputum at rest, nighttime, or after minor exercise; trouble sleeping or waking up at night or using more pillows to prop yourself at night because of breathing problems; weight gain; or foot or leg swelling.

Call your doctor right away if you notice that your heartbeat or pulse is irregular. This is a sign that you are in AFib.

MULTAQ may cause liver problems, including life-threatening liver failure. Tell your doctor right away if you are taking MULTAQ and develop any of these signs and symptoms of liver problems: loss of appetite; nausea; vomiting; fever; feeling unwell; unusual tiredness; itching; yellowing of the skin or the whites of the eyes; unusual darkening of the urine; right upper stomach area pain or discomfort.

An additional serious side effect is a slowed heartbeat.

You should not take MULTAQ if you have severe liver problems or if you had liver problems after using amiodarone.

For Medication Guide and full Prescribing Information, including boxed WARNING, please refer to the links on MULTAQ.com, the site from which this page was downloaded.

Please see additional Important Safety Information on next page.

Important Safety Information for MULTAQ® (dronedarone) (continued)

You should not take MULTAQ if you are breastfeeding, pregnant, or plan to become pregnant as it is not known if it may pass to or harm your baby. Women who may become pregnant should use effective birth control while taking MULTAQ. Talk to your doctor about the best birth control methods for you.

MULTAQ can interact with certain medications. Do not use the following medications with MULTAQ:

- Nefazodone
- Voriconazole
- Ritonavir
- Telithromycin
- Ketoconazole
- Clarithromycin
- Itraconazole
- Cyclosporine

The following medicines can lead to a dangerous abnormal heart rhythm if taken with MULTAQ:

- Phenothiazines
- Tricyclic antidepressants
- Certain antibiotics
- Certain medicines used to treat an abnormal heart rhythm or fast heartbeat

Your potassium levels should be within the normal range prior to and while taking MULTAQ. Tell your doctor about any diuretic medications (“water pills”) you are taking as some can lower the level of potassium or magnesium in your blood.

The most common side effects when taking MULTAQ include stomach problems such as diarrhea, nausea, vomiting, stomach area pain, and indigestion; feeling tired and weak; skin problems such as redness, rash, and itching. Tell your doctor about any side effect that bothers you or that does not go away.

Do not drink grapefruit juice while taking MULTAQ, as this can increase the likelihood that you will have a side effect of MULTAQ.

Tell your doctor about all medications, vitamins, and herbal remedies (including St. John’s Wort) you take to help avoid serious drug interactions.

Your doctor will monitor your heart rhythm regularly to make sure your heartbeat keeps a normal rhythm. Your doctor may perform certain tests, such as heart exams and blood tests, while you’re taking MULTAQ.

Please see additional Important Safety Information on previous page.

If you’ve been prescribed MULTAQ, make sure to ask your doctor these important questions:

- Question: What is severe heart failure? What should I do if I start to experience symptoms of worsening heart failure? (Some of these symptoms include shortness of breath or wheezing at rest; wheezing, chest tightness, or coughing up frothy sputum at rest, nighttime, or after minor exercise; trouble sleeping or waking up at night or using more pillows to prop yourself at night because of breathing problems; weight gain; or foot or leg swelling.)

Notes: _____

- Question: What are some of the symptoms of liver problems and what should I do if I experience them? (Some of these symptoms include: loss of appetite; nausea; vomiting; fever; feeling unwell; unusual tiredness; itching; yellowing of the skin or the whites of the eyes; unusual darkening of the urine; right upper stomach area pain or discomfort.)

Notes: _____

- Question: What should I know about pregnancy and taking MULTAQ? Can I take MULTAQ while breast feeding?

Notes: _____

- Question: What should I do if I experience a serious side effect, like slowed heartbeat, or other side effects, like diarrhea, nausea, or vomiting?

Notes: _____

- Question: What should I do if I have AFib symptoms or notice that my pulse is irregular while I’m taking MULTAQ?

Notes: _____

- Question: I understand that you will need to monitor my heart rhythm regularly while I’m taking MULTAQ to make sure my heartbeat keeps a normal rhythm. How often should I come in? Can I schedule those visits now?

Notes: _____

- Question: What information do I need to share with my other doctors now that I’m taking MULTAQ?

Notes: _____

For Medication Guide and full Prescribing Information, including boxed WARNING, please refer to the links on MULTAQ.com, the site from which this page was downloaded.

