MULTAQ® (dronedarone) 400mg Tablets

GETTING STARTED with MULTAQ

Your guide to resources and support



INDICATION

MULTAQ is a prescription medicine used to lower the chance of hospitalization for atrial fibrillation (AFib) in people who currently have a normal heart rhythm and have had certain types of AFib (paroxysmal or persistent AFib) in the past. It is not known if MULTAQ is safe and effective in children younger than age 18 years old.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about MULTAQ?

MULTAQ may cause serious side effects, including:

- **Increased risk of death, stroke, and heart failure in people with:**
  - **A certain type of heart failure called decompensated heart failure.** Heart failure is when your heart does not pump blood through your body as well as it should. MULTAQ can cause new or worsening heart failure.
    Do not take MULTAQ if you have symptoms of heart failure that recently worsened and you were hospitalized, or if you have severe heart failure.
    Call your healthcare provider right away if you develop any of the following signs or symptoms of heart failure during treatment with MULTAQ: shortness of breath or wheezing at rest; wheezing, chest tightness, or coughing up frothy sputum at rest, nighttime, or after minor exercise; trouble sleeping or waking up at night because of breathing problems; using more pillows to prop yourself up at night so you can breathe more easily; gaining more than 5 pounds quickly; increasing swelling of feet or legs

Please see additional Important Safety Information throughout, and read the **Medication Guide** and full **Prescribing Information including Risk of SERIOUS SIDE EFFECTS**.

MULTAQ® (dronedarone) 400mg Tablets



STEPS TO GETTING STARTED

So, your doctor has told you that you have paroxysmal or persistent AFib—and while your heart is currently in normal rhythm, you have been prescribed MULTAQ. What do you do next?

Follow these steps to help you start your treatment!

STEP 1

Know how to take MULTAQ

Take MULTAQ exactly as your doctor tells you and read the helpful information in Step 1 about how to take it.

STEP 2

Get help paying for MULTAQ

Learn if you qualify for savings support options.

STEP 3

Keep track of your next visit and questions for your doctor

Be sure to write down questions you have for your doctor to help with your conversation.

Take some time to review this information along with a loved one or caregiver so they can help support you in your treatment with MULTAQ.

IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I should know about MULTAQ? (cont'd)

- **A certain type of irregular heartbeat (rhythm) called permanent atrial fibrillation (AFib).** Permanent AFib is when you and your healthcare provider decide not to try to change your heart rhythm back to a normal heart rhythm or your heart rhythm cannot be changed back to a normal rhythm.
- **Do not take MULTAQ** if you have permanent AFib. Your healthcare provider should check your heart rhythm regularly to make sure your heart keeps a normal rhythm.
- Call your healthcare provider right away if you develop any of the following signs or symptoms of AFib during treatment with MULTAQ such as: fast or irregular heartbeat or pulse; chest pain; dizziness or lightheadedness; tiredness or weakness; reduced ability to exercise; shortness of breath

Please see additional Important Safety Information throughout, and read the **Medication Guide** and full **Prescribing Information including Risk of SERIOUS SIDE EFFECTS**.

STEP 1

KNOW HOW TO TAKE MULTAQ®

Take MULTAQ as prescribed



Pill not actual size

For MULTAQ to work the way it is supposed to, you must take it exactly as your doctor tells you. Ask your doctor any questions you have about how to take MULTAQ.

Remember, do not stop taking MULTAQ without first talking with your doctor.

Take 1 tablet twice a day with food, once with your morning meal and once with your evening meal

Be sure your meal includes a healthy fat. A healthy fat may help the body better absorb the medicine. Examples of healthy fats are avocados, olive oil, and unsalted nuts.*

If you miss a dose of MULTAQ, skip the missed dose and take your next dose at your regular time

Do not take 2 doses at the same time to make up for a missed dose.

If you take too much MULTAQ, call your doctor or go to the nearest hospital emergency room right away

Do not drink grapefruit juice during treatment with MULTAQ

Grapefruit juice can increase the amount of MULTAQ in your blood. This, in turn, can increase the chances of getting side effects.

DRUG INTERACTIONS

MULTAQ and certain other medicines can interact with each other, causing serious side effects. Let your doctors know about all medications you are currently taking (including vitamins and herbal remedies).

Visit MULTAQ.com
for tips on taking MULTAQ.

*Avoid nuts if a nut allergy exists. Talk to your doctor about a diet that is right for you.

Sanofi does not provide medical advice, diagnosis, or treatment. The health information contained herein is provided for general educational purposes only. Your healthcare professional is the best source of information regarding your health. Please consult your healthcare professional if you have any questions about your health or treatment.

IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I should know about MULTAQ? (cont'd)

MULTAQ doubles your risk of dying if you have these conditions. Your healthcare provider may give you a medicine to help prevent blood clots and decrease your risk of stroke during treatment with MULTAQ. Tell your healthcare provider right away if you develop any of the following signs or symptoms of stroke during treatment with MULTAQ such as: numbness or weakness in the face, arms, or legs, especially on 1 side of the body; confusion, trouble speaking, or difficulty understanding things; trouble seeing in 1 or both eyes; trouble walking, dizziness, loss of balance, or lack of coordination.

Please see additional Important Safety Information throughout, and read the Medication Guide and full Prescribing Information including Risk of SERIOUS SIDE EFFECTS.

STEP 2

GET HELP PAYING FOR MULTAQ®

Over 90% of commercially insured patients paid $0 per fill with use of the MULTAQ Savings Card*



You may be eligible to **pay as low as $0 copay** on each of your **MULTAQ fills,** with no income restrictions, using the MULTAQ Savings Card.† Terms and conditions apply.

Visit MULTAQ.com to see if you are eligible.

*Based on 2022-2023 claims data.

†**MULTAQ Savings Program: This savings program is not insurance.**

This offer is not valid for prescriptions covered by or submitted for reimbursement, in whole or in part, under Medicare, Medicaid, VA, DOD, TRICARE, similar federal or state programs, including any state pharmaceutical programs. If you have an Affordable Care (Health Care Exchange) plan, you may still be qualified to receive and use this savings card. Please note: the Federal Employees Health Benefits (FEHB) Program is not a federal or state government health care program for purposes of the savings program. Void where prohibited by law. For the duration of the program, eligible commercially insured patients may be eligible for a [$0] co-pay per monthly fill. Cash patients may be eligible for up to [$150] off per monthly fill. Maximums apply. Savings may vary depending on patients' out-of-pocket costs. The MULTAQ Savings Program applies to the cost of medication. There are other relevant costs associated with overall treatment. Sanofi reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Upon registration, patients will receive all program details. For questions regarding your eligibility or benefits, or if you wish to discontinue your participation, call the MULTAQ Savings Program at (833) 950-1304 (9:00 am-7:00 pm EST, Monday-Friday).

IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I should know about MULTAQ? (cont'd)

- **Liver problems.** MULTAQ may cause severe liver problems, including life-threatening liver failure. **Do not take MULTAQ** if you have severe liver problems. Your healthcare provider may order blood tests to check your liver before you start taking MULTAQ and during treatment.

Call your healthcare provider right away if you develop any of the following signs and symptoms of liver problems during treatment with MULTAQ: loss of appetite, nausea, vomiting; fever, feeling unwell, unusual tiredness; itching; yellowing of the skin or the whites of the eyes (jaundice); unusual darkening of the urine; right upper stomach area pain or discomfort

Please see additional Important Safety Information throughout, and read the **Medication Guide** and full **Prescribing Information including Risk of SERIOUS SIDE EFFECTS**.

GET HELP PAYING FOR MULTAQ®

For people with low income or Medicare

Sanofi Patient Connection®

This patient assistance program provides MULTAQ to people with limited income and resources and who also meet other qualifications.

PATIENT Connection®

Visit sanofipatientconnection.com to download an application and see if you qualify.

MEDICARE EXTRA HELP

If you qualify, you may be able to pay no more than $10.35 monthly through Extra Help from Medicare.

Visit Medicare.gov to learn more.

IMPORTANT SAFETY INFORMATION (CONT'D)

Who should not take MULTAQ?

See **"What is the most important information I should know about taking MULTAQ?"**
Do not take MULTAQ if:

- you have a certain type of heart problem called heart block, and you do not have an implanted pacemaker
- your heart rate is less than 50 beats each minute
- you have had liver or lung problems after using amiodarone
- you have a certain type of electrocardiogram (ECG) abnormality including QTc or PR interval prolongation

Please see additional Important Safety Information throughout, and read the **Medication Guide** and full **Prescribing Information including Risk of SERIOUS SIDE EFFECTS**.

STEP 3

KEEP TRACK OF YOUR QUESTIONS AND VISITS

IMPORTANT SAFETY INFORMATION (CONT'D)

Who should not take MULTAQ? (cont'd)

See **"What is the most important information I should know about taking MULTAQ?"**

Do not take MULTAQ if:

- you take certain medicines that can change the amount of MULTAQ that gets into your body such as: nefazodone; ritonavir; ketoconazole; itraconazole; erythromycin; voriconazole; telithromycin; clarithromycin; cyclosporin
- you take certain medicines that can lead to a dangerous abnormal heart rhythm such as: phenothiazines; tricyclic antidepressants; macrolide antibiotics; certain medicines for abnormal heart rhythm or fast heartbeat (Class I and III antiarrhythmics)
- you are allergic to dronedarone or any of the other ingredients in MULTAQ

What should I tell my healthcare provider before taking MULTAQ?

Before taking MULTAQ, tell your healthcare provider about all of your medical conditions, including if you:

- have any other heart problems, including heart rhythm problems, or have had a stroke
- have an implanted pacemaker
- have liver or kidney problems
- have lung problems
- have low levels of potassium or magnesium in your blood
- are pregnant or plan to become pregnant. MULTAQ may harm your unborn baby

  Females who can become pregnant
  - Your healthcare provider will do a pregnancy test before you start treatment with MULTAQ
  - Use effective birth control (contraception) during treatment and for 5 days after your final dose of MULTAQ
  - Tell your healthcare provider right away if you think you are pregnant or become pregnant during treatment with MULTAQ
- are breastfeeding or plan to breastfeed. It is not known if MULTAQ passes into your breast milk. Do not breastfeed during treatment and for 5 days after the final dose of MULTAQ

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking MULTAQ with certain other medicines may affect the amount of MULTAQ or other medicines in your blood and may increase your risk of side effects or affect how well MULTAQ or the other medicines work.

Especially tell your healthcare provider if you take: medicine for high blood pressure, chest pain, or other heart conditions; statin medicine to lower blood cholesterol; medicine for tuberculosis (TB); medicine for seizures; digoxin; warfarin or other blood thinner medicines; medicine for organ transplant; an herbal supplement called St. John's wort; water pills (diuretics).

Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.

What should I avoid while taking MULTAQ?

Do not drink grapefruit juice during treatment with MULTAQ. Grapefruit juice can increase the amount of MULTAQ in your blood and can increase your chance of getting side effects.

What are the possible side effects of MULTAQ?

MULTAQ may cause serious side effects, including:

- See **"What is the most important information I should know about MULTAQ?"**
- **Inflammation of the lungs, including scarring and thickening.** Call your healthcare provider if you develop shortness of breath or a dry cough during treatment with MULTAQ
- **Low potassium and magnesium levels in your blood.** This can happen if you take certain water pills (diuretics) during treatment with MULTAQ. Your healthcare provider may check you for this problem before and during treatment. Tell your healthcare provider if you develop any of the following symptoms of low potassium or low magnesium during treatment with MULTAQ: nausea or vomiting; weakness or sleepiness; muscle weakness, spasms, or tremors; loss of appetite; constipation; heart palpitations; tingling or numbness

MULTAQ® (dronedarone) 400mg Tablets



Find support and savings at **MULTAQ.com/ways-to-save**.

ON MULTAQ.COM, YOU CAN:

- **Learn more about AFib**
- **Review how to take MULTAQ** (2X A DAY)
- **Find programs to help pay for MULTAQ**

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of MULTAQ? (cont'd)

- **Changes in the electrical activity in your heart called QT interval prolongation.** QT interval prolongation can increase your chance of getting dangerous abnormal heart rhythms
- **Kidney problems and kidney failure.** MULTAQ can cause changes in kidney function that can be serious and lead to kidney failure, especially in people with heart failure or people with low body fluid levels. Your healthcare provider will check your blood for signs of kidney problems during treatment. Tell your healthcare provider if you develop any of the following symptoms of kidney problems during treatment with MULTAQ: loss of appetite; nausea and vomiting; muscle cramps; dry, itchy skin; swelling of the feet and ankles; shortness of breath; trouble sleeping; urinating too much or too little

The most common side effects of MULTAQ include: diarrhea; weakness, lack of energy, and feeling very tired or sleepy (asthenia); nausea; skin problems such as redness, rash, and itching; stomach area (abdominal) pain; slow heart rate (bradycardia); vomiting; indigestion.

Your healthcare provider may stop treatment with MULTAQ if you develop certain side effects. These are not all of the possible side effects of MULTAQ.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see additional Important Safety Information throughout, and read the **Medication Guide** and full **Prescribing Information including Risk of SERIOUS SIDE EFFECTS**.

If you are a patient experiencing problems with a Sanofi U.S. product, please contact Sanofi U.S. at 1-800-633-1610.

© 2026 Sanofi. All rights reserved. MULTAQ is a registered trademark of Sanofi or an affiliate. MAT-US-2402872-v2.0-03/2026

sanofi