

## MULTAQ® (dronedarone) Health Care Professional Checklist

This checklist is intended to assist health care professionals with identifying the appropriate patients for MULTAQ. MULTAQ is indicated to reduce the risk of hospitalization for atrial fibrillation (AFib) in patients in sinus rhythm with a history of paroxysmal or persistent AFib. MULTAQ is available in 400-mg tablets.

### The following are contraindications for use with MULTAQ.

**Permanent AFib** that will not or cannot be cardioverted into normal sinus rhythm

**Symptomatic heart failure with recent decompensation requiring hospitalization or NYHA Class IV heart failure**

2nd or 3rd degree AV block, or sick sinus syndrome (except when used in conjunction with a functioning pacemaker)

Bradycardia <50 bpm

Concomitant use of strong CYP 3A inhibitors, such as ketoconazole, itraconazole, voriconazole, cyclosporine, telithromycin, clarithromycin, nefazodone, and ritonavir

Concomitant use of drugs or herbal products that prolong the QT interval and might increase the risk of Torsade de Pointes, such as phenothiazine antipsychotics, tricyclic antidepressants, certain oral macrolide antibiotics, and Class I and III antiarrhythmics

Liver or lung toxicity related to the previous use of amiodarone

A QTc Bazett interval  $\geq 500$  ms or PR interval  $> 280$  ms

Severe hepatic impairment

Pregnancy (Category X): patients who are or may become pregnant

Nursing mothers

Hypersensitivity to the active substance or to any of the excipients

### The following information was derived from WARNINGS and PRECAUTIONS

<ul style="list-style-type: none"><li>• Cardiac rhythm should be monitored (<math>\leq 3</math> months)</li></ul>	<ul style="list-style-type: none"><li>• Potassium levels should be within normal range prior to and maintained in the normal range during administration of MULTAQ</li></ul>
<ul style="list-style-type: none"><li>• MULTAQ should only be initiated in patients in sinus rhythm who are receiving appropriate antithrombotic therapy</li></ul>	<ul style="list-style-type: none"><li>• Onset of dyspnea or non-productive cough may be related to pulmonary toxicity and patients should be carefully evaluated clinically</li></ul>
<ul style="list-style-type: none"><li>• Advise patients to consult a physician if they develop signs or symptoms of heart failure</li></ul>	<ul style="list-style-type: none"><li>• Monitor renal function periodically</li></ul>
<ul style="list-style-type: none"><li>• Consider obtaining periodic hepatic serum enzymes, especially during the first 6 months of treatment</li></ul>	

Please see Important Safety Information on next page, and for full Prescribing Information, including boxed WARNING, please [click here](#) or visit [MULTAQ.com/hcp](http://MULTAQ.com/hcp).

# Important Safety Information for MULTAQ® (dronedarone)

## **WARNING: INCREASED RISK OF DEATH, STROKE AND HEART FAILURE IN PATIENTS WITH DECOMPENSATED HEART FAILURE OR PERMANENT ATRIAL FIBRILLATION**

**MULTAQ is contraindicated in patients with symptomatic heart failure with recent decompensation requiring hospitalization or NYHA Class IV heart failure. MULTAQ doubles the risk of death in these patients.**

**MULTAQ is contraindicated in patients in atrial fibrillation (AFib) who will not or cannot be cardioverted into normal sinus rhythm. In patients with permanent AFib, MULTAQ doubles the risk of death, stroke, and hospitalization for heart failure.**

### **MULTAQ is also contraindicated in patients:**

- With second- or third-degree atrioventricular (AV) block or sick sinus syndrome (except when used in conjunction with a functioning pacemaker), bradycardia <50 bpm, QTc Bazett interval  $\geq$ 500 ms or PR interval >280 ms
- Who are or may become pregnant (Category X) or nursing. MULTAQ may cause fetal harm when administered to a pregnant woman
- With concomitant use of strong CYP 3A inhibitors, such as ketoconazole, itraconazole, voriconazole, cyclosporine, telithromycin, clarithromycin, nefazodone, ritonavir, or drugs or herbal products that prolong the QT interval and might increase the risk of Torsade de Pointes, such as phenothiazine antipsychotics, tricyclic antidepressants, certain oral macrolide antibiotics, and Class I and III antiarrhythmics
- With liver or lung toxicity related to the previous use of amiodarone
- With severe hepatic impairment
- With hypersensitivity to the active substance or to any of the excipients

### **Cardiovascular Death in NYHA Class IV or Decompensated Heart Failure**

MULTAQ is contraindicated in patients with NYHA Class IV heart failure or symptomatic heart failure with recent decompensation requiring hospitalization because it doubles the risk of death.

### **Cardiovascular Death and Heart Failure in Permanent AFib**

MULTAQ doubles the risk of cardiovascular death (largely arrhythmic) and heart failure events in patients with permanent AFib. Patients treated with MULTAQ should undergo monitoring of cardiac rhythm no less often than every 3 months. Cardiovert patients who are in AFib (if clinically indicated) or discontinue MULTAQ. MULTAQ offers no benefit in subjects in permanent AFib.

### **Increased Risk of Stroke in Permanent AFib**

In a placebo-controlled study in patients with permanent AFib, dronedarone was associated with an increased risk of stroke, particularly in the first two weeks of therapy. MULTAQ should only be initiated in patients in sinus rhythm who are receiving appropriate antithrombotic therapy.

### **New Onset or Worsening Heart Failure**

New onset or worsening of heart failure has been reported during treatment with MULTAQ in the postmarketing setting. In a placebo-controlled study in patients with permanent AFib, increased rates of heart failure were observed in patients with normal left ventricular function and no history of symptomatic heart failure, as well as those with a history of heart failure or left ventricular dysfunction.

Advise patients to consult a physician if they develop signs or symptoms of heart failure, such as weight gain, dependent edema, or increasing shortness of breath. If heart failure develops or worsens and requires hospitalization, discontinue MULTAQ.

### **Liver Injury**

Hepatocellular liver injury, including acute liver failure requiring transplant, has been reported in patients treated with MULTAQ in the postmarketing setting. Advise patients treated with MULTAQ to report immediately symptoms suggesting hepatic injury (such as anorexia, nausea, vomiting, fever, malaise, fatigue, right upper quadrant pain, jaundice, dark urine, or itching). Consider obtaining periodic hepatic serum enzymes, especially during the first 6 months of treatment. It is not known whether routine periodic monitoring of serum enzymes will prevent the development of severe liver injury. If hepatic injury is suspected, promptly discontinue MULTAQ and test serum enzymes, aspartate aminotransferase (AST), alanine aminotransferase (ALT), and alkaline phosphatase, as well as serum bilirubin, to establish whether there is liver injury. If liver injury is found, institute appropriate treatment and investigate the probable cause. Do not restart MULTAQ in patients without another explanation for the observed liver injury.

### **Pulmonary Toxicity**

Cases of interstitial lung disease including pneumonitis and pulmonary

fibrosis have been reported in patients treated with MULTAQ in the post-marketing setting. Onset of dyspnea or non-productive cough may be related to pulmonary toxicity and patients should be carefully evaluated clinically. If pulmonary toxicity is confirmed, MULTAQ should be discontinued.

### **Hypokalemia and Hypomagnesemia with Potassium-Depleting Diuretics**

Hypokalemia and hypomagnesemia may occur with concomitant administration of potassium-depleting diuretics. Potassium levels should be within the normal range prior to administration of MULTAQ and maintained in the normal range during administration of MULTAQ.

### **QT Interval Prolongation**

MULTAQ induces a moderate (average of about 10 ms but much greater effects have been observed) QTc (Bazett) prolongation. If the QTc Bazett interval is  $\geq$ 500 ms, discontinue MULTAQ.

### **Renal Impairment and Failure**

Marked increase in serum creatinine, pre-renal azotemia and acute renal failure, often in the setting of heart failure or hypovolemia, have been reported in patients taking MULTAQ. In most cases, these effects appear to be reversible upon drug discontinuation and with appropriate medical treatment. Monitor renal function periodically.

Small increases in creatinine levels (about 0.1 mg/dL) following MULTAQ treatment initiation have been shown to be a result of inhibition of creatinine's tubular secretion. The elevation has a rapid onset, reaches a plateau after 7 days and is reversible after discontinuation.

### **Women of Childbearing Potential**

Pre-menopausal women who have not undergone a hysterectomy or oophorectomy must use effective contraception while using MULTAQ. Dronedarone caused fetal harm in animal studies at doses equivalent to recommended human doses. Counsel women of childbearing potential regarding appropriate contraceptive choices.

### **Drug-Drug Interactions**

- Treatment with Class I or III antiarrhythmics or drugs that are strong inhibitors of CYP 3A must be stopped before starting MULTAQ (see Contraindications)
- Patients should be instructed to avoid grapefruit juice beverages while taking MULTAQ
- Calcium channel blockers with depressant effects and beta-blockers could increase the bradycardia effects of MULTAQ on conduction
- In the ANDROMEDA (patients with recently decompensated heart failure) and PALLAS (patients with permanent AFib) trials, baseline use of digoxin was associated with an increased risk of arrhythmic or sudden death in MULTAQ-treated patients compared to placebo. In patients not taking digoxin, no difference in risk of sudden death was observed in the MULTAQ vs placebo groups
- Digoxin can potentiate the electrophysiologic effects of MULTAQ (such as decreased AV-node conduction). MULTAQ increases exposure to digoxin. Consider discontinuing digoxin. If digoxin treatment is continued, halve the dose of digoxin, monitor serum levels closely, and observe for toxicity
- Postmarketing cases of increased INR with or without bleeding events have been reported in warfarin-treated patients initiated with MULTAQ. Monitor INR after initiating MULTAQ in patients taking warfarin
- Statins: Avoid simvastatin doses greater than 10 mg daily. Follow statin label recommendations for use with CYP 3A and P-gP inhibitors such as MULTAQ

### **Adverse Reactions**

In studies, the most common adverse reactions observed with MULTAQ were diarrhea, nausea, abdominal pain, vomiting, and asthenia.

**For full Prescribing Information, including boxed WARNING, please [click here](#) or visit [MULTAQ.com/hcp](http://MULTAQ.com/hcp).**